

PROJECTED FUTURE CONDITIONS IN THE LOWER WILLAMETTE RIVER SUBBASIN OF NORTHWEST OREGON: CLACKAMAS, MULTNOMAH & WASHINGTON COUNTIES

Roger Hamilton, Bob Doppelt, Steve Adams & Stacy Vynne
(University of Oregon, Climate Leadership Initiative)

Maps and Graphs by Kathie Dello and Darrin Sharp
(Oregon Climate Change Research Institute)

With Support from the Lower Willamette Advisory Team

December 2009



OREGON CLIMATE CHANGE RESEARCH INSTITUTE



CLIMATE LEADERSHIP INITIATIVE
INSTITUTE FOR A SUSTAINABLE ENVIRONMENT
UNIVERSITY OF OREGON

TABLE OF CONTENTS

INTRODUCTION	3
MODELS & LIMITATIONS	4
GLOBAL CLIMATE CHANGE PROJECTIONS	6
SELECT RESOURCES FOR CURRENT ENVIRONMENTAL CONDITIONS WITHIN THE LOWER WILLAMETTE SUBBASIN	7
CLIMATE PROJECTIONS FOR THE LOWER WILLAMETTE: CLACKAMAS, MULTNOMAH AND WASHINGTON COUNTIES OF NORTHWEST OREGON	8
TEMPERATURE	8
PRECIPITATION	10
VEGETATION	11
SNOW WATER EQUIVALENT	13
STREAMFLOW	13
ACRES BURNED	14
APPENDIX A: SPATIAL MAPS AND GRAPHS OF MODELING RESULTS	17

INTRODUCTION

The Lower Willamette River Subbasin of western Oregon is rich in history, culture, and biological diversity. For the purposes of this assessment we have defined the Subbasin as the counties of Clackamas, Multnomah, and Washington, including the mouth of the Willamette River (flowing into the Columbia River) and the Tualatin and Clackamas river subbasins.

The Lower Willamette Subbasin is the most downstream (yet northern) portion of the Willamette River Basin. It is the most extensively urbanized Subbasin in this assessment with 51% in urban land use. Most of the greater Portland metropolitan region is in this Subbasin, with 25 cities, the urban portions of three counties and about 40 percent of the state's population. Portland, home to 570,000 people, is the largest city in the state with 2.2 million people in the greater metropolitan area.

The entire Willamette Basin stretches from Cottage Grove in the south to Portland in the north. The Basin is nestled in between the Cascade Range to the East and the Coastal Range to the West. Major tributaries include the Calapooya, Clackamas, Coast Fork, Long Tom, Luckiamute, McKenzie, Mary's, Middle Fork, Molalla, Santiam, Tualatin and Yamhill Rivers. With its fertile soils, the River Basin has attracted settlers since the early 1800s. The entire Willamette Basin includes a land area of 11,500 square miles, a

population of 2.5 million and about 75% of Oregon's economy. While the Lower Basin is less than a third of the landmass in the entire River Basin, it contains the majority of the Basin and state's population and economy.



Figure 1. Willamette River Basin (in yellow) with focus for this report in red (Washington, Multnomah & Clackamas counties)

This report is intended to provide an ecological overview of the Subbasin and localized projections of the consequences of climate change in the Lower Willamette Subbasin. It is provided to support climate preparedness and adaptation planning and policy development in the Lower Willamette. The climate change models presented in this report were mapped by scientists at the Oregon Climate Change Research Institute. The Climate Leadership Initiative at the University of Oregon helped develop this summary of the assessment.

MODELS & LIMITATIONS

Preparing and planning for climate change is, above all, an exercise in risk management. Traditionally, future planning has been based on historical conditions and experiences. However, this approach is no longer reliable as climate change will produce never before seen changes in temperature, precipitation, streamflow, vegetation and fire patterns. To understand the possible impacts on natural, built, economic, human and cultural systems, climate models are used to project future conditions.

Understanding what actions should be taken to prepare natural, built, human, cultural and economic systems for climate change is challenging as the Earth's climate and ocean systems are too complex to be simulated in a laboratory experiment or reactor. Therefore, climate scientists use global climate models to estimate how climate change might affect conditions in mid- and end-of-century. Climate models incorporate the physical laws and chemical interactions of the Earth. Future conditions are calculated based on different "scenarios" (or estimations) of future greenhouse gas emissions, policies and regulations that would limit emissions, technological improvements, and behavioral changes. (For the scenarios selected in this project, please see below.) In order to test the climate models, they are backcasted against observed data to see how well they "predict" the past. While each of the inputs to the models are the same, they vary in their level of detail and manner of

interpretation. This results in some differences in outputs and uncertainty as to which future scenario is most likely to occur (and therefore the importance of running multiple models). The difference in detail and interpretation that leads to this uncertainty is due to processes and feedbacks between different parts of the Earth's climate system that are not fully understood. However, by comparing a group of climate models, it is possible to project a credible range of possible future conditions.

Most climate models are created at global scales, but are difficult to downscale to local or regional scales because the more localized they become the greater the chance of errors and uncertainty. However, managers and policymakers need regional and local data that reflect how climate change will impact their region in order to plan and develop policies. The Oregon Climate Change Research Institute (OCCRI) has adjusted global model results to local and regional scales to support this effort.

The Intergovernmental Panel on Climate Change (IPCC) uses approximately 27 models to make global climate projections. The models are developed by different institutions in different countries around the world and are subject to different interpretations.

OCCRI has selected the following models for use in the Lower Willamette Basin project:

- **PCM1:** The Parallel Climate

Model, developed through a collaboration of United States federal agencies.

- **CSIRO-MK3**: Developed by the Atmospheric Research Office in Australia.
- **HadCM3**: Developed by the Met Office, the national weather office for the United Kingdom.
- **MIROC**: A Japanese model used for the MC1 vegetation models (shown in results for fire and vegetation projections).

These models were selected because they use temperature and precipitation forcing agents including changes in greenhouse gas emissions, aerosols, water vapor and cloud cover, solar radiation, and changes in land use to represent possible future conditions.

To further refine these projected futures, the IPCC has developed a range of scenarios under which climate models are run. These scenarios, as described in the IPCC's Special Report on Emissions Scenarios (SRES), describe different futures for greenhouse gas emissions, land use, and agricultural practices based on global policy decision-making.¹ For this report, two scenarios were selected to model how different futures might play out:

- **A1b**: The business as usual scenario (for which current global emissions are actually exceeding) that presumes continued growth in economies, population and technology, and reliance on mixed energy sources.
- **B1**: The 'greener' emissions scenario, which suggests emissions increasing slightly in the coming decades but then

falling to lower than current levels by 2100 due to deployment of low carbon energy and transportation systems.

Model outputs were converted to local scales using local data on recent temperature and precipitation patterns. The MC1 vegetation model provides information on possible future vegetation types and wildfire patterns.

The utility of the model results presented in this report is to assist public and private entities envision what the conditions and landscape may look like in the future and the potential magnitude and direction of change.

It is important to note that the scenarios described in this report utilize the best available information. However, they are not predictions. Instead, they should be considered **possible outcomes**. Actual conditions may vary quite substantially from those depicted in these scenarios. Readers are therefore urged to **focus on the range of projections and the trends they suggest**, as opposed to relying on the outputs of a single model or on a particular number

¹ For more information on SRES, visit:
http://www.ipcc-data.org/ddc_envdata.html

GLOBAL CLIMATE CHANGE PROJECTIONS

The IPCC² and the U.S. Global Change Research Program³ agree that the evidence is “unequivocal” that the Earth’s atmosphere and oceans are warming, and that this warming is due primarily to human activities including the emission of CO₂, methane, and other greenhouse gases, along with land conversion and deforestation. Average global air temperature has already increased by 0.7° C (1.4° F) over the last hundred years and is expected to increase up to 6.4° C (11.5° F) within the next century (Figure 1).

Even with immediate reductions in greenhouse gas emissions, impacts from the current build up of greenhouse gas emissions in the atmosphere will continue to be felt for decades. It may take decades or centuries to restabilize the system. Reducing emissions is a vital mitigation measure to prevent further impacts on climate systems. However, countries and communities must also begin to plan and prepare for the likely impacts that will be experienced as a result of the emissions already present in the

² IPCC 2007. Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.

³ USGCRP 2009. Global Climate Change Impacts in the United States. T. R. Karl, J. M. Melillo, and T. C. Peterson, (eds.). Cambridge University Press.

atmosphere. By taking proactive steps to plan for changes, residents of the Lower Willamette will be better positioned to build resistance and resiliency of the systems they depend on for maintaining quality of life under a climate changed future.

When using projections to prepare for climate change, we must consider how to deal with the uncertainty of models and make decisions that are robust against a range of future scenarios. One approach is finding consistency in models; if consistencies do not exist, then we must consider strategies that are effective no matter what change occurs. This is likely to involve building system resiliency and resistance, as well as flexibility into the planning process.

Global Warming Projections




Figure 2. Projections for global temperature increase for a number of models used by the IPCC, compared with temperatures over the last one hundred years. Note that while projections for temperature increase vary by the end of the century, all models show a clear upward trend. (From IPCC 2007)

SELECT RESOURCES FOR CURRENT ENVIRONMENTAL CONDITIONS WITHIN THE LOWER WILLAMETTE SUBBASIN

City of Portland Endangered Species Program
<http://www.portlandonline.com/bes/fish/index.cfm>

City of Portland Invasive Plants Strategy Report. 2008
<http://www.portlandonline.com/bes/index.cfm?c=47815&a=217069>

City of Portland - Willamette Watershed Characterization Report
<http://www.portlandonline.com/BES/index.cfm?c=31806>

Oregon Department of Environmental Quality. 2008 Air Annual Report.
<http://www.deq.state.or.us/aq/forms/2008AQreport.pdf>

Department of Environmental Quality Willamette Basin Report. 2009.
<http://www.deq.state.or.us/about/eqc/agendas/attachments/2009oct/F-WillametteBasinAssessmentRpt.pdf>

Oregon State University - Institute for Natural Resources - Oregon Explorer
http://inr.oregonstate.edu/oregon_explorer.html

Oregon State University - Oregon Natural Heritage Information Center
<http://oregonstate.edu/ornhic/index.html>

Oregon State University - Willamette Basin Explorer
<http://www.willametteexplorer.info/>

CLIMATE PROJECTIONS FOR THE LOWER WILLAMETTE: CLACKAMAS, MULTNOMAH AND WASHINGTON COUNTIES OF NORTHWEST OREGON

Outputs of our climate models (PCM, CSIRO, and HadCM) and the vegetation model (MC1) include projections for changes in temperature, precipitation, percent of landscape burned, suitable vegetation types and distribution, snowpack, and streamflow. A historical baseline of 1971-2000 was used in order to make comparisons of projections for the 2040s (1930-1959) and 2080s (2070-2099) (scientists use thirty year time slices, or averages, to account for interannual and interdecadal variability). Stream data is for 2020s and 2040s, due to data availability. The results present a range of different possible future conditions in the Lower Willamette. Unforeseen circumstances such as uncertainties about chemical reactions or international policy to drastically reduce greenhouse gas emissions, may result in a future different than has been projected.

Climate change projections are provided here as bar graphs, charts and spatial maps to demonstrate the results of the modeling using a variety of visualizations




Figure 3. Sample of spatial maps for summer temperature projections. PCM1 Model projections for both scenarios with historic baseline (1971-2000), mid- and end-century projections. Darker red indicates warmer temperatures.

that may be useful for different decision-making groups. Samples for each factor are posted below: the full suite of maps, charts and graphs can be found at the end of the document.

The modeling results come from the global modeling results available from the IPCC Fourth Assessment Report. Implications for the Pacific Northwest are based on the twenty global climate models analyzed by Mote and Salathé (2009). For historical baseline, 800m PRISM 1971-2000 climate grids were used to apply to the analysis and 'downscale' the data.

Temperature

The overall trend for temperature shows warming for the entire Lower Willamette by the end of the century. The most intense

warming of 10-15° F is in summer. Warming is also most intense along the Interstate 5 corridor (running through the most populated area of the Basin). There is also warming during the winter months, but less extreme than in the summer (about 3-5° F). In the B1 scenario, there is still warming, but warming is less severe in the summer months compared with the higher emissions scenario, and minimal in winter.




Figure 4. Projections for seasonal change in temperature for all models and both scenarios.

Precipitation

Precipitation is one of the variables most difficult for climate models to project, particularly for the Pacific Northwest. For the mid-century A1b scenario, an increase in precipitation is seen in the winter for all models (minimal for PCM1), with least change in the spring, and mixed results in the summer. HadCM shows severe drought in the summer, while CSIRO and PCM1 show a slight increase in precipitation. For the fall months, the models project mixed results in precipitation. In the 2080s, the CSIRO and PCM1 models show an increase in precipitation for most of the year. HadCM projects slight reduction in the spring and drought in the summer, but not as severe a drought as shown in the 2040s.




Figure 5. Projections for percent change in precipitation compared to historical (1971-2000) for all models under the A1b scenario.






Figure 6. Sample spatial maps for summer projections for the HadCM model for both scenarios. Darker red indicates less precipitation.

Vegetation

For Maritime Evergreen Needleleaf species, the HadCM and MIROC models project a significant decline, with near disappearance by the end of the century. CSIRO also shows a decline, but not as severe as the other models. HadCM shows a rapid increase in Temperate Evergreen Needleleaf species, replacing Maritime species. MIROC and CSIRO also project an increase in Temperate Evergreen species, but not as rapidly as HadCM. Subtropical Mixed Forest species also increases slightly in abundance, with rapid increase after mid-century under the MIROC model, but less change for HadCM and CSIRO. (Data provided by Ray Drapek, Pacific Northwest Research Station.)



Figures 7-8. Changes in dominant vegetation for both scenarios using the MC1 model.

Snow Water Equivalent

Under the A1b scenario, the model projects a severe decrease in snow water equivalent with near disappearance (greater than 80% loss) by the end of the century. (Data provided by Heejun Chang, Portland State University.)




Figure 9. Ensemble mean changes in April 1 snow water equivalent for A1b scenario.

Streamflow

Projections show that streams are likely to become flashier in the winter and early spring– that is, higher high flows and more frequent and severe flooding, and lower low flows with more streams going dry – due to temperature, more precipitation falling as rain, groundwater and storm severity changes. All three models show a slight increase in winter flow, with moderate decrease in historical summer flows.




Figure 10. Sample stream flow change for 2020s and 2040s under the HadCM model.

Acres Burned

The data below shows percent of area in each grid cell (8km) projected to be burned for the entire Willamette Basin (note, counties for the Lower Subbasin could not be overlaid). Under both scenarios, HadCM projects a greater proportion burned with almost 2% of each grid cell burned by 2080. MIROC and HadCM show an increase in intensity of areas burned, especially under the A1b scenario and for the Lower Willamette Subbasin. CSIRO, which tends to be a wetter and cooler model, shows less change in proportion burned. (Data provided by Ray Drapek, Pacific Northwest Research Station.)




Figure 11. Percent area burned for all models under both scenarios.

Appendix A: Spatial Maps and Graphs of Modeling Results

A1-2 Biomass consumed by fire for HadCM, MIROC and CSIRO

A3-4 Proportion biomass burned for HadCM, MIROC and CSIRO

A5-6 Vegetation carbon for HadCM, MIROC and CSIRO


A7-8 Change in vegetation type for HadCM, MIROC and CSIRO

A9-20. Seasonal temperature spatial maps for HadCM, CSIRO, PCM1


A21-32 Seasonal precipitation spatial maps for HadCM, CSIRO, PCM1

A33-52 Streamflow graphs for HadCM, MIROC and CSIRO


Willamette Basin Vegetation Carbon Consumed By Fire: A1B Emissions




Willamette Basin Vegetation Carbon Consumed By Fire: B1 Emissions




Willamette Basin Proportion Of Vegetation Consumed By Fire: A1B Emissions




Willamette Basin Proportion Of Vegetation Consumed By Fire: B1 Emissions





Willamette Basin Vegetation Carbon: A1B Emissions



Willamette Basin Vegetation Carbon: B1 Emissions





Predominant Vegetation Type

b1

a1b

CSIRO3.5

HadCM3







MIROC3.2

CSIRO3.5







HadCM3

MIROC3.2







1971-2000



2040s



2080s




-  Subalpine Forest
-  Maritime Evergreen Needleleaf Forest
-  Temperate Evergreen Needleleaf Forest
-  Temperate Evergreen Needleleaf Woodland
-  Temperate Grassland
-  Subtropical Mixed Forest




Fall

CSIRO3.5




Fall

HadCM3



Fall

PCM1



Spring

CSIRO3.5




March

April




May

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s




Temperature (F)



Spring

HadCM3



Spring

PCM1




March

April




May

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s



Temperature (F)



Summer

CSIRO3.5




June

July




August

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s



Temperature (F)



Summer

HadCM3




June

July




August

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s



Temperature (F)



Summer

PCM1




June

July




August

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s



Temperature (F)



Winter

CSIRO3.5




December

January




February

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s




Temperature (F)




Winter

HadCM3




Temperature (F)




Winter

CSIRO3.5




Temperature (F)




Fall

CSIRO3.5




Precipitation (in.)



Fall

HadCM3



Fall




PCM1

September

October




November

1971-2000






b1




2040s



2080s






1971-2000






a1b


2040s



2080s




Precipitation (in.)




Spring

HadCM3



Precipitation (in.)



Winter

CSIRO3.5




December

January




February

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s




Precipitation (in.)




Spring

HadCM3




Winter

HadCM3




Spring

PCM1



Precipitation (in.)



Winter

PCM1




December

January




February

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s



Precipitation (in.)



Summer

CSIRO3.5




June

July




August

b1




1971-2000



2040s






2080s






a1b




1971-2000




2040s



2080s




Precipitation (in.)




Summer

HadCM3




Summer


PCM1



October	12089.0518	10625.6875	8551.72754
November	43828.4453	49754.2188	46570.9297
December	67830.4922	78045.9531	77284.2578
January	69016.1719	71506.375	71506.375
February	63242.8203	66327.3125	66327.3125
March	52075.6211	57860.7617	57860.7617
April	43373.5156	44029.4766	44029.4766
May	31571.2148	27501.6797	27501.6797
June	21786.75	15417.5176	15417.5176
July	10500.6689	6036.96143	6036.96143
August	3766.96899	1851.12805	1851.12805
September	3490.72217	2249.32617	2249.32617



Willamette River above Falls at Oregon City




— Historical

— 2020s

— 2040s




Willamette River at Portland




HadCM3

Clackamas River above Three Lynx Creek




Molalla River near Canby




HadCM3

Clackamas River at River Mill Dam



HadCM3


Tualatin River at West Linn



— Historical — 2020s — 2040s




Willamette River above Falls at Oregon City




— Historical — 2020s — 2040s




Willamette River at Portland




Clackamas River above Three Lynx Creek




Molalla River near Canby




Clackamas River at River Mill Dam



Tualatin River at West Linn




Willamette River above Falls at Oregon City




PCM1

Willamette River at Portland




PCM1

Clackamas River above Three Lynx Creek




Molalla River near Canby




PCM1

Clackamas River at River Mill Dam



Tualatin River at West Linn



— Historical

— 2020s

— 2040s

